




Fulvic START



PANATOP FULVIC START

- **Innovative mix with two main purposes:**

- › For seed treatment
- › Foliar application in the earliest phase of the crop growth
- › The balanced ratio of fulvic and humic acids, amino acids, phosphorus (P) and the high content of zinc (Zn), manganese (Mg) and copper (Cu) have a strong stimulating effect on the seed germination and the beginning of tillering.

- **Humic acids contribute to:**

- › The transformation of nutrients into a form available for the plants
- › They absorb moisture and stimulate germination.

- **The combination of phosphorus (P) and highly concentrated zinc (Zn), manganese (Mn) and copper (Cu) stimulate:**

- › Rooting and tillering in the most important growth stage of the crops.
- › The result is uniform sprouting, well-developed root system, increased health status and preservation of the yield potential at the time of its formation.

Panatop Fulvic Start is the budget solution for every farm.

<h3>Fulvic START®</h3> <p>www.panamin.bg</p>																						
<table border="1"> <thead> <tr> <th colspan="2">COMPOSITION</th> </tr> </thead> <tbody> <tr> <td>Phosphorus pentoxide (PO₅)</td> <td>5,80%</td> </tr> <tr> <td>L-amino acids</td> <td>5,80%</td> </tr> <tr> <td>Fulvic and humic acids</td> <td>2,32%</td> </tr> <tr> <td>Organic matter in dry matter</td> <td>70,41%</td> </tr> <tr> <td>Organic carbon</td> <td>3,48%</td> </tr> </tbody> </table>		COMPOSITION		Phosphorus pentoxide (PO ₅)	5,80%	L-amino acids	5,80%	Fulvic and humic acids	2,32%	Organic matter in dry matter	70,41%	Organic carbon	3,48%									
COMPOSITION																						
Phosphorus pentoxide (PO ₅)	5,80%																					
L-amino acids	5,80%																					
Fulvic and humic acids	2,32%																					
Organic matter in dry matter	70,41%																					
Organic carbon	3,48%																					
<table border="1"> <thead> <tr> <th colspan="2">TRACE ELEMENTS</th> </tr> </thead> <tbody> <tr> <td>Zinc (Zn) EDTA</td> <td>0,58%</td> </tr> <tr> <td>Manganese (Mn) EDTA</td> <td>0,58%</td> </tr> <tr> <td>Copper (Cu) EDTA</td> <td>0,23%</td> </tr> <tr> <td>Molybdenum (Mo) EDTA</td> <td>0,06%</td> </tr> </tbody> </table>		TRACE ELEMENTS		Zinc (Zn) EDTA	0,58%	Manganese (Mn) EDTA	0,58%	Copper (Cu) EDTA	0,23%	Molybdenum (Mo) EDTA	0,06%											
TRACE ELEMENTS																						
Zinc (Zn) EDTA	0,58%																					
Manganese (Mn) EDTA	0,58%																					
Copper (Cu) EDTA	0,23%																					
Molybdenum (Mo) EDTA	0,06%																					
<table border="1"> <thead> <tr> <th colspan="2">PARAMETER</th> </tr> </thead> <tbody> <tr> <td>Density</td> <td>1,12 + 0,2 % kg/L</td> </tr> <tr> <td>pH value</td> <td>6,5</td> </tr> </tbody> </table>		PARAMETER		Density	1,12 + 0,2 % kg/L	pH value	6,5															
PARAMETER																						
Density	1,12 + 0,2 % kg/L																					
pH value	6,5																					
<table border="1"> <thead> <tr> <th colspan="2">APPLICATION</th> </tr> </thead> <tbody> <tr> <td>Foliar</td> <td>1-1,5 L/Ha</td> </tr> <tr> <td>Soil treatment</td> <td>6-8 L/Ha</td> </tr> <tr> <td>Winter wheat – from the beginning of tillering until the forming of rosette</td> <td>every 10-14 days</td> </tr> <tr> <td>Spring wheat – from 3-5th leaf until 2-3 leaf pair</td> <td>every 10-14 days</td> </tr> <tr> <td>Perennial crops – beginning: after blooming</td> <td>every 10-14 days</td> </tr> <tr> <td>Vegetables – beginning: before, after transplanting at 10-15 cm height</td> <td>every 10-14 days</td> </tr> <tr> <td rowspan="3">Seed treatment</td> <td>Cereals</td> <td>1,5-2 L/1000 kg seeds</td> </tr> <tr> <td>Corn</td> <td>5 L/1000 kg seeds</td> </tr> <tr> <td>Sundowner</td> <td>7,5 L/1000 kg seeds</td> </tr> </tbody> </table>		APPLICATION		Foliar	1-1,5 L/Ha	Soil treatment	6-8 L/Ha	Winter wheat – from the beginning of tillering until the forming of rosette	every 10-14 days	Spring wheat – from 3-5th leaf until 2-3 leaf pair	every 10-14 days	Perennial crops – beginning: after blooming	every 10-14 days	Vegetables – beginning: before, after transplanting at 10-15 cm height	every 10-14 days	Seed treatment	Cereals	1,5-2 L/1000 kg seeds	Corn	5 L/1000 kg seeds	Sundowner	7,5 L/1000 kg seeds
APPLICATION																						
Foliar	1-1,5 L/Ha																					
Soil treatment	6-8 L/Ha																					
Winter wheat – from the beginning of tillering until the forming of rosette	every 10-14 days																					
Spring wheat – from 3-5th leaf until 2-3 leaf pair	every 10-14 days																					
Perennial crops – beginning: after blooming	every 10-14 days																					
Vegetables – beginning: before, after transplanting at 10-15 cm height	every 10-14 days																					
Seed treatment	Cereals	1,5-2 L/1000 kg seeds																				
	Corn	5 L/1000 kg seeds																				
	Sundowner	7,5 L/1000 kg seeds																				

PANATECHNOLOGY RECOMMENDATIONS:

**Panatop Fulvic Start 0,5-1,5 L/Ha + Panamin Suspension 1-2 L/Ha
or Panatop Alga Max 0,5-1 L/Ha; Panatop BLOOM 1-1,5 L/Ha;
Panatop Amino Zn 1-1,5 L/Ha**